



Cambodia • Lao PDR • Thailand • Viet Nam
For sustainable development


BioRA DSS Workshop

Preliminary Calibration: Geomorphology

BioRA DSS Technical Workshop
Phnom Penh, Cambodia
15-19 February 2016

www.mrcmekong.org

Contents



- Calibration scenarios
- Geomorphology results by indicator
 - Erosion
 - Bed material grain-size
 - Availability exposed sandy habitat
 - Availability inundated sandy habitat
 - Availability exposed rocky habitat
 - Availability inundated rocky habitats
 - Depth of bedrock pools
 - Water clarity
- Final comments

www.mrcmekong.org

Calibration Scenarios




Scenario	Characteristics
CS1	High dry season flow, low wet season flow
CS2	6 dry years, followed by 6 wet years, etc.
CS3	A shortened wet season
CS4	Sediment supply at 75% of Preliminary Reference
CS5	Migration blocked between FA1 and FA2 ONLY
CS7	Extreme dry year (1992 – 10%) repeated for whole sequence
CS8	Migration blocked between FA4 and 5 ONLY
CS9	Migration blocked between FA1 and 2 <u>AND</u> between FA4 and 5
CS10	Sediment supply at 25% of Preliminary Reference

www.mrcmekong.org

Comments on Calib Scenarios



- Flow & sediments are typically correlated in the LMB
- Each calib scenarios only changes **one** parameter
 - Either flow or sediment
 - Needs to be recognised in interpretation
- Migration 'blocks' do not affect sediment or flow
 - Not Applicable to geomorph
- Underlying trends in sediment data sets



www.mrcmekong.org

Changes to Indicators

Scenarios CS5, CS8, CS9 Not applicable to Geomorphology



Indicators	Calibration scenarios									
	CS1	CS2	CS3	CS4	CS5	CS7	CS8	CS9	CS10	
Discipline : Geomorphology										
Erosion (bank / bed incision)	-4.9	0.7	-10.1	9.2	0.7	-8.7	0.7	0.7	36.4	
Average bed sediment size - dry season	1.0	0.8	0.6	1.2	0.3	1.1	0.3	0.3	2.5	
Availability exposed sandy habitat - dry season	7.9	-5.3	4.1	-4.5	0.2	4.8	0.2	0.2	-20.8	
Availability inundated sandy habitat -dry season	1.9	0.4	2.3	-5.2	-2.0	-0.4	-2.0	-2.0	-15.4	
Availability exposed rocky habitat -dry season	-1.0	-5.0	-7.3	7.2	1.0	-3.3	1.0	1.0	25.2	
Availability inundated rocky habitat -dry season	<0.1	5.8	-3.9	5.4	0.1	-5.2	0.1	0.1	20.6	
Depth of bedrock pools in dry season	-7.1	-0.3	-4.7	5.3	0.2	-7.7	0.2	0.2	12.3	
Water clarity	1.2	27.5	38.9	16.4	1.2	53.6	1.2	1.2	242.4	

www.mrcmekong.org

Changes to Indicators

■ <-20
 ■ 20 - 50%
 ■ 50 - 70%
 ■ >70%




Indicators	Calibration scenarios									
	CS1	CS2	CS3	CS4	CS5	CS7	CS8	CS9	CS10	
Discipline : Geomorphology										
Erosion (bank / bed incision)	-4.9	0.7	-10.1	9.2		-8.7			36.4	
Average bed sediment size - dry season	1.0	0.8	0.6	1.2		1.1			2.5	
Availability exposed sandy habitat - dry season	7.9	-5.3	4.1	-4.5		4.8			-20.8	
Availability inundated sandy habitat -dry season	1.9	0.4	2.3	-5.2		-0.4			-15.4	
Availability exposed rocky habitat -dry season	-1.0	-5.0	-7.3	7.2		-3.3			25.2	
Availability inundated rocky habitat -dry season	<0.01	5.8	-3.9	5.4		-5.2			20.6	
Depth of bedrock pools in dry season	-7.1	-0.3	-4.7	5.3		-7.7			12.3	
Water clarity	1.2	27.5	38.9	16.4		53.6			242.4	

Erosion All Seasons

Response Curves based on:
Shear stress \propto Flow
 $Flow_{wet} \gg Flow_{dry}$

Sediment $\propto 1/Erosion$
 $\downarrow Sediment_{Load} = \uparrow Erosion$




CS	Characteristics	%Change	Rationale
CS1	High dry flow, low wet flow	-4.9	Dec in wet flow reduces erosion more than inc in dry season increases erosion
CS2	6 dry, 6 wet	0.7	Inc. in dry & dec in wet = net '0'
CS3	A shortened wet season	-10	Reduced erosion from reduced flows
CS4	Sediment supply at 75%	+9	Inc due to less sediment avail for transport
CS7	Extreme dry year	-9	Low flow = red shear stress with same sed
CS10	Sediment supply at 25%	+36	Inc due to less sediment avail for transport

www.mrcmekong.org

Erosion at FA1: CS1

- High dry season, low wet season
- High shear stress in dry, low in wet
- High sediment load in dry
- Dec in erosion due to reduced wet combined with inc. sediment load in dry
- Change = -5%




www.mrcmekong.org



Erosion at FA1: CS2


- 6 dry, 6 wet years
- Lower in dry years
- Higher in wet years
- T1 sed & shear stress v. low
- No Net change
 - Change = <1%



www.mrcmekong.org

Erosion at FA1: CS3:

- Short wet season
 - Wet duration is lower (103 v 143)
- Sed duration lower
 - 54 days v 62 days
- Shear stress the same as P Ref
- Change = -10%




www.mrcmekong.org

Erosion at FA1: CS4 & CS10



- Sediment reduction
 - CS4: Sed load = 75%
 - CS10: Sed load = 25%
- All other parameters the same
- Change
 - CS4: Erosion increases by 9%
 - CS10: Erosion increases by 36%




www.mrcmekong.org



Erosion at FA1: CS7



- Extreme dry year
- V. low sediment concentrations
- Reduced shear stress
- Change:
 - Erosion decreases by 10%




www.mrcmekong.org



Sandy Habitat at FA1: CS1


- High Dry, Low Wet
- Decrease in erosion
 - Erosion CS1= -5%
- Max dry season WL
 - Increase by 0.3 m
- High sed load in dry
- Dec in erosion greater influence than increase in water level
- Net change +8%



www.mrcmekong.org

Sandy Habitat at FA1: CS2


- 6 Dry, 6 Wet
- No net change in erosion
- Max dry season WL
 - Increase by 0.55 m
- Inc during dry years
- Decrease during wet
- Net change +8%



www.mrcmekong.org

Sandy Habitat at FA1: CS3


- Short Wet Season
- Decrease in erosion
 - -10%
- Max dry season WL
 - Increase by 0.6 m
- Inc during dry years
- Small net change +4%



www.mrcmekong.org

Sandy Habitat at FA1: CS4 & CS10


- Reduction in sediment
 - CS4: 75% P Ref
 - CS10: 25% P Ref
- Water levels the same
 - Difference is due to erosion
- Increase in erosion results in less availability
 - CS4: -5% availability
 - CS10: -20% availability



www.mrcmekong.org

Sandy Habitat at FA1: CS7


- Extreme dry
- Lower water levels
- Dec in erosion
- 5% Increase in availability



www.mrcmekong.org

Avail Inundated Sandy Habitat

Dependant on:
Water level
Erosion




CS	Characteristics	%Change	Rationale
CS1	High dry flow, low wet flow	2	Changes on an annual scale but virtually no change – increases & decreases balance out over the period of record
CS2	6 dry, 6 wet	<1	
CS3	A shortened wet season	2	
CS4	Sediment supply at 75%	-5	Reduced sediment reduces availability of sandy habitat
CS7	Extreme dry year	<-1	Low water levels decrease availability
CS10	Sediment supply at 25%	-15	Loss of sandy habitat due to increased erosion

www.mrcmekong.org

Sandy Habitat at FA1: Compare Exposed & Inundated CS10




- 25% Sediment load
- Large increase in erosion
- Decrease in both exposed & inundated availability



www.mrcmekong.org

Avail Exposed Rocky Habitat- Dry Season

Dependant on:
Water level
Erosion




CS	Characteristics	%Change	Rationale
CS1	High dry flow, low wet flow	-1	Annual changes but no net change
CS2	6 dry, 6 wet	-5	Dry increases, wet decreases & net is slight decrease
CS3	A shortened wet season	-7	Dec in erosion covers rock with sediment
CS4	Sediment supply at 75%	7	Reduced sediment increases erosion so more rock exposed
CS7	Extreme dry year	-3	Increased deposition reduces rock availability
CS10	Sediment supply at 25%	25	Large increase in erosion exposes more rock

www.mrcmekong.org



Sandy Habitat at FA1: Compare Exposed Sandy & Exposed Rocky CS10


- 25% Sediment load
- Large increase in erosion
- Inc in rocky habitat as availability of sandy decreases



www.mrcmekong.org

Avail Inundated Rocky Habitat- Dry Season

Dependant on:
Water level
Erosion




CS	Characteristics	%Change	Rationale
CS1	High dry flow, low wet flow	<1	Change in annual pattern but no net change
CS2	6 dry, 6 wet	6	Dry decreases, wet increases & net is slight increase
CS3	A shortened wet season	-4	Dec in erosion covers rock with sediment
CS4	Sediment supply at 75%	5	Increase in erosion increases exposure of rock
CS7	Extreme dry year	-5	Lower water levels decrease availability
CS10	Sediment supply at 25%	20.6	Large increase in erosion exposes more rock

www.mrcmekong.org

Depth of Bedrock Pools Dry Season


Dependant on:
Erosion
Sediment timing
Water level




CS	Characteristics	%Change	Rationale
CS1	High dry flow, low wet flow	-7	Low wet reduces scour of pools
CS2	6 dry, 6 wet	<-1	Large annual changes but no net change
CS3	A shortened wet season	-5	Reduced erosion to scour pool
CS4	Sediment supply at 75%	5	Less sediment results in less deposition
CS7	Extreme dry year	-8	Reduced erosion to scour pool
CS10	Sediment supply at 25%	12	Less sediment results in less deposition so depth increases

www.mrcmekong.org

Water Clarity



Dependant on:
Sediment Load
Grain-size



CS	Characteristics	%Change	Rationale
CS1	High dry flow, low wet flow	1	No change to sediment loads
CS2	6 dry, 6 wet	28	V. Large increase during dry seasons
CS3	A shortened wet season	39	Increased duration of low sed conc.
CS4	Sediment supply at 75%	16	Red. Sediment increases clarity
CS7	Extreme dry year	54	Dry year has low sediment concentrations
CS10	Sediment supply at 25%	242	V. low sediment concentrations

www.mrcmekong.org

Final comments

- Calibration scenarios results are consistent with physical processes & Mekong characteristics
- Small changes each year
 - Combine to create change
 - Balance out in 'dynamic equilibrium'
 - E.g. 6-dry, 6 wet have distinct annual signal but create little change overall
- Final calibration after DSF results available

www.mrcmekong.org

Cambodia • Lao PDR • Thailand • Viet Nam
For sustainable development



Thank You

www.mrcmekong.org